

Research Article

Extraction of Total Lignans from Radix Isatidis and Its Ant-RSV Virus Effect

Hui-qin XU¹, Li-Wei HE^{1,2*}, Ying-yue DU², Shu Zhang²

¹Department of Pharmacology, School of Pharmacy, Nanjing University of Chinese Medicine, Nanjing 210023, China

²Department of Pharmacology, Nanjing University of Chinese Medicine Hanlin College, Taizhou, 225312, China

Abstract

The orthogonal test method was used to investigate the effects of solvent volume fraction, extraction time and solvent on the extraction process. The results showed that the best extraction process of the total lignans in Radix Isatidis is 70% ethanol, the extraction time is 2 h, and the amount of solvent is 8 times. The inhibitory effect of the total lignans of Radix Isatidis on RSV was determined by MTT method and cytopathic effect (CPE) method. The total lignans of Radix Isatidis significantly reduced the formation of plaques during the phase of virus adsorption and virus replication. The plaques decreased slightly when the virus entered the cell stage, indicating that the total lignans of Radix Isatidis may exert their resistance during the stage of virus adsorption and virus replication. The role of the virus. The extraction conditions of the Radix lignans obtained by orthogonal design experiments are stable and feasible.

Keywords: Radix isatidis; Total LIGNANS; Orthogonal design; Antiviral

Introduction

Radix Isatidis is a traditional Chinese medicine, derived from the dry roots of the cruciferous plant *Isatis indigotica*. It has the effect of clearing heat and detoxifying cold blood and pharynx [1,2]. Nowadays, Radix Isatidis is often used clinically for the treatment of bacterial infections and viral diseases, including influenza virus, herpes simplex virus, respiratory syncytial virus (RSV), mumps virus, hepatitis B virus, coxsackie virus and so on [3-6]. It is commonly considered a broad-spectrum natural antiviral drug [7]. However, due to the complex chemical composition of Radix Isatidis, the discovery of the potential medicinal components of Radix Isatidis and the mechanism of action is seriously restricted [8]. The view that there may be a synergistic effect of Banlangen on the human body has been reported. The theory of "pharmacy superposition" proposed by Professor Cai [9] of Peking University shows that most of the active ingredients with similar active structures act on one or more similar targets. Different types of compounds have been isolated from Radix isatidis including indirubin, clemastanin B, alkaloids, lignans, and flavonoids [10,11]. The chemical composition of Banlangen mainly includes three major categories: alkaloids, lignans and organic acids. At present, studies on Banlangen focus on the study of total alkaloids

and total organic acid sites [12-14], and there are few reports on the study of lignans. However, the antiviral effect of lignans is significant [15,16], and the total lignans in Radix Isatidis occupies a larger proportion, so the antiviral effect of total lignans in Banlangen cannot be ignored. The extraction process of total lignans from Radix Isatidis was optimized, and its antiviral effect was studied. It provides a theoretical and practical basis for the effective utilization of Radix Isatidis medicinal resources and the synergistic antiviral effects of Radix Isatidis.

Materials

Cells and Viruses

Human laryngeal carcinoma epithelial cells (Hep2 cells) were kindly provided by the Institute of Pediatrics, Nanjing University of Chinese Medicine; RSV type A Long strain was provided by Wuhan Institute of Virology, Chinese Academy of Sciences.

Drugs and Reagents

Banlangen (Origin: Bozhou, Anhui Province, batch number: 20171211), medicinal herbs were provided by the market of Anhui Bozhou medicinal herbs, were identified to meet the "People's Republic of China Pharmacopoeia" (2015 edition) standards; sodium hydroxide (batch number:150708); hydrochloric acid (batch number: 20141126); Methanol (analytical alcohol, batch number: 20170110); Ethanol (batch number: 20161216); Ribavirin (Shanghai Yuanye Bio, Batch No. 36791045); DMEM high-glucose medium (US HyClone, Lot No. AAK207235); Pancreatin, Fetal calf serum (US GIBCO, Lot 1801539, 20171026); MTT, DMSO, BSA (Sigma USA, batch number C10036912, 045M4080V, 10735078001); RIPA, phenylmethylsulfonyl fluoride, BCA kit, loading Buffer (Shanghai Biyuntian Biological Co., Ltd., batch number is 0123251105, 0421323721, 03161712517, 0140260519); Phosphatase inhibitor, protease inhibitor (Jiangsu Jikai Biological Technology Co., Ltd., batch number KGP602180214, KGP603024523); anti-GAPDH antibody, goat anti-rabbit fluorescent secondary antibody; Protein Marker (Runsheng Biological Co., Ltd.,

Citation: Hui-qin XU, Li-Wei HE, Ying-yue DU, Shu Zhang. Extraction of Total Lignans from Radix Isatidis and Its Ant-RSV Virus Effect. *Clin J Pharmacol Pharmacother*. 2019; 1(1): 1005.

Copyright: © 2019 Hui-qin XU

Publisher Name: Medtext Publications LLC

Manuscript compiled: Feb 13th, 2019

***Corresponding author:** Liwei He, Department of Pharmacology, Nanjing University of Chinese Medicine Hanlin College, P. O. Box: 225312, No. 6 Xueyuan Road, Taizhou, China, Tel: +86 0523 80639253; Fax: +86 0523 80639253; E-mail: hlw9752@163.com

batch number 03018120); RSV F antibody (batch number sc-101362, purchased from Santa Cruz Biotechnology); RSV G antibody (batch ab94966, purchased from Santa Cruz Biotechnology); 1:29 bis-Acr, 10% SDS, APS (US GIBCO, Lot No. B0153K0103185, 3642A462, B016M052548); TRIS, Glycine (Beijing Soley Company, Lot 562B018, 803B0156); Skim milk powder (Shanghai Wulian Technology Co., Ltd., Lot No. 1216542).

Methods and Results

Optimization of total lignan extraction process of Radix Isatidis

Using L_9 (3^4) orthogonal design (Table 1) [17,18], the volume fraction, solvent amount and extraction time of ethanol were investigated, and the extraction rate and total lignans content obtained under various conditions were determined. The total lignans amount was measured by ultraviolet spectrophotometry. Using larch resin alcohol as a reference substance, through full-wavelength scanning, it can be seen that the reference substance and test solution have the same maximum absorption wavelength at 280nm, so it is determined to measure at 280nm. Pipette precision pipette a certain volume of larch resin alcohol standard solution, place it in a 10ml volumetric flask, add methanol to dilute to volume, and shake, to obtain concentrations of 0.01mg/ml, 0.02mg/ml, 0.04mg/ml, 0.08 mg/ml, 0.16 mg/ml, 0.32 mg/ml standard solution, respectively. The methanol solution was used as a blank solution to determine the absorbance at the maximum absorption wavelength of the above prepared standard solutions with different concentrations. The concentration of the standard solution was plotted on the abscissa, and the absorbance value A was plotted on the ordinate to draw a standard curve. The total lignans content was calculated and the results are shown in Table 2.

Orthogonal test results show that the total rate of lignans in Radix Isatidis decreases with increasing ethanol volume fraction. According to the extreme difference of the Extraction rate, the effects of various factors on the rate of total lignans were as follows: ethanol volume fraction > extraction time > solvent amount. The pharmacological effect was determined by the total lignans content of Radix Isatidis. Therefore, the total lignan content is the main indicator in this experiment. Because the amount of solvent has little effect on the experimental results, adhering to the principle of saving resources, 8 times the amount of solvent is selected for extraction. Therefore, the preferred process conditions are 70% ethanol volume fraction, 2 h extraction time, and 8 times solvent Table 3.

Macroporous resin enrichment and purification

The total lignan content of Radix Isatidis was extracted according to the preferred process conditions in "2.1", the solvent was recovered in a low pressure environment, and the sample was prepared to have a mass concentration of 1 g/ml, adjusted to pH=7 with NaOH, and the macroporous resin [19,20] was reprocessed and then wet-packed. Column, the sample was applied to the column, statically adsorbed for 12 hours, and eluted with distilled water to remove unadsorbed components and water-soluble impurities, and then

Table 1: Extraction process factor level.

Level	A ethanol volume fraction/%	B extraction time/h	C solvent amount
1	70	1	6
2	80	1.5	8
3	95	2	12

eluted with different volume fractions of ethanol solution, identified by ultraviolet spectrophotometry, and identified as the eluent of the lignan component was dried under reduced pressure to obtain the total lignan of Radix isatidis [21,22].

Determination of total lignan content

Accurately refer to the "2.2" item the total lignan sample 0.6mg, placed in a 10ml volumetric flask, diluted with methanol to determine the volume, that is, the sample solution. The total lignan content was determined under the same conditions as in "2.1" item. The total lignan content in the sample after isolation and purification was 74.87%. Total lignan determination methodological investigation.

Precision Inspection

A 1 ml of the test solution was taken and measured according to the method of "2.1" (n=6), and the result showed that the RSD value was 0.235%. It shows that the precision of the instrument is good.

Reproducibility Study

Six samples (n=6) of the total lignan sample of Radix Isatidis were taken in parallel and measured according to the method of "2.1". The results showed that the RSD value was 2.03%, indicating that the method has good reproducibility.

Stability Investigation

Take the appropriate amount of total lignans from Radix Isatidis, dissolve with methanol, and make up to volume. Take 1 ml of the test solution, according to the method of 2.1, and measure the absorbances at the time of 0 min, 5 min, 15 min, 30 min, 1 h and 2 h, the absorbance was 0.2305, 0.2386, 0.2317, 0.2400, 0.2339 and 0.1538, respectively. It indicates that the sample solution has good stability within 1 h.

Recovery Test

The sample was accurately weighed, and an appropriate amount of the reference substance was added and measured according to the above method. The recovery rates of the samples were calculated to be 98.59%, 103.15%, 98.76%, 99.53%, 102.58%, and 98.36%, respectively.

Antiviral effects of total lignans from Radix Isatidis

Cell lines and Virus: Hep2 cells were cultured in DMEM high glucose medium with 10% FBS, and virus amplification and virus virulence assay methods were described in 21. Calculate virus TCID₅₀=10^{-5.5}/50ul according to Reed-Muench formula.

Cell proliferation assay: Hep2 cells were seeded at 1×10⁵ in 96-well plates at 200 μL per well. DMEM (without FBS) was used to dialysis the drug to 7 different concentrations. When the cells are full of the bottom of the holes, the above-mentioned formulated drugs were added. Continue to culture in a constant temperature incubator for 24 h, then add 10 ul of 5% MTT 22 per well, incubate in the incubator for 4 hours, remove the culture solution, add 100 ul of DMSO per well, shake at low speed for 10 min, and measure the absorbance at 490 nm on a microplate reader. The results are shown in Figure 1 and 2. The results showed that the concentration of lignans in Radix Isatidis was less than 200 μg/ml had no effect on cell survival, and that higher than 200 μg/ml all inhibited the proliferation of Hep2 cells in a dose-dependent manner, so the maximum non-toxicity of total lignans in Radix Isatidis The concentration is about 200 μg/ml. As shown in Figure 2, the concentration of ribavirin below 500 μg/ml had no effect on cell survival, and higher than 500 μg/ml dose-dependently inhibited the proliferation of Hep2 cells, so the maximum nontoxic concentration of ribavirin was about 500 μg/ml.

Table 2: orthogonal test scheme and results of extraction process.

Test sequence	A ethanol volume fraction/%	B extraction time/h	C solvent amount	D	Total lignan content/%	Extract rate/%
1	1	1	1	1	-	24.63
2	1	2	2	2	-	24.89
3	1	3	3	3	-	25.05
4	2	1	2	3	0.07	23.75
5	2	2	3	1	0.09	23.24
6	2	3	1	2	0.06	23.46
7	3	1	3	2	0.08	23.91
8	3	2	1	3	0.07	22.45
9	3	3	2	1	0.06	22.82
Total Lignan content K1	0	0.15	0.13	0.15		
K2	0.22	0.16	0.13	0.14		
K3	0.21	0.12	0.16	0.14		
R	0.09	0.04	0.03	0.01		
Extract Rate K1	74.57	72.29	70.54	70.69		
K2	70.45	70.58	71.46	72.26		
K3	69.18	71.33	72.20	71.25		
R	5.39	1.71	1.66	1.57		

Table 3: Analysis of variance.

Factors	Sum of squares of deviations	df	F	P
A	8.823	2	46.44	>0.05
B	0.309	2	1.724	>0.05
C	0.693	2	3.869	>0.05
D	0.346	2		

Direct anti-RSV study of total lignans

Hep2 cells were seeded in 6-well plates, and different concentrations of total lignans were incubated with RSV on ice. The whole process is carried out on ice to prevent the virus from entering the cell. After 2 h of incubation, the cells were washed with pre-cooled medium to remove unbound virus. The cells were then lysed and the RSV G and F proteins used to detect cell attachment were analyzed by immunoblot [23]. The result is shown in Figure 3. Hep2 cells were seeded in 6-well plates and incubated with RSV (500 µL virus titer of 100 TCID50/well) for 2 hours on ice, allowing virus attachment in the presence or absence of the indicated concentration of drug. The non-attached virus was removed with ice-cold medium and then allowed to adsorb at 37°C for 2 hours to allow the attached RSV to enter the host cell. At the end of the incubation period, the culture medium was removed and the culture was replaced with agarose containing 1% of 2.5% calf serum and incubated at 37°C for 6-7 days. The formation of plaques was observed daily. The result is shown in Figure 4.




Figure 1: The effect of lignans of Radix Isatidis on Hep2 cells survival rate (x±s, n=6).

Hep2 cells were inoculated into 6-well plates, 500 ul of 100 TCID50 virus solution was added to each well, and placed in a constant temperature incubator for 2 h, shaking once every 20 min. After the end of the adsorption, the virus suspension in the culture plate was discarded, and replaced with agarose containing 2.5% calf serum 1%, and different concentrations of the drug were added, and the mixture was incubated at 37 °C to observe the formation of plaques every day. The result is shown in Figure 4.




Figure 2: The effect of Ribavirin on Hep2 cell survival rate (x±s, n=6).

The RSV F protein and RSV G protein were not expressed in the normal cell group, and the RSV F protein and RSV G protein expression levels were higher in the virus group, indicating that the virus adsorption amount was higher in 2 hours. Compared with the virus group, each drug group RSV The expression of F protein and RSV G protein decreased to varying degrees (P<0.01). With the increase of drug concentration, the expression of RSV F protein and RSV G protein decreased in turn.

As shown in Figure 3, the addition of lignans only slightly reduced the number of plaques at the stage of entry of the virus into the cells ,

indicating that the compound does not exert its antiviral effect against RSV entering the cell stage. At the virus binding and replication stage, as shown in Figure 3 (black bars and red bars), the drugs in the total lignans of Radix is added, causing a significant reduction in plaques at this stage, and the number of plaques decreased with the increase of concentration of lignans, it indicates that the compound may exert its antiviral effect by hindering virus adsorption and inhibiting viral replication. In the stage of virus replication, total lignans inhibit viral replication less than ribavirin group.

Discussion

The chemical composition of Radix Isatidis is very complicated, which seriously hinders the research process of Radix Isatidis. It has been shown in the literature that compounds with larger content may play a leading role in the overall content and efficacy of such components [9], and the total lignan content has a large proportion in Banlangen, and its antiviral effect is diverse and the mechanism of action is not clear. The study on the antiviral effect of total lignans is of progressive significance to elucidate the antiviral mechanism of Radix Isatidis, so its antiviral effect needs to be proved. However, since the lignans are mainly polar substances and there is no specific color reaction and identification method, it is difficult to enrich, because the research in this part is still blank, so the extraction and purification of the total lignans from Radix Isatidis the process needs to be unified.

In this study, we used orthogonal test to optimize the extraction process of total lignans from Radix Isatidis. The effects of ethanol concentration, solvent dosage, and extraction time on the total lignans content of Isatis indica were investigated. The method is stable, reliable and suitable for applicability; macroporous resin adsorption separation technique was used to separate and enrich the extracted samples. For the problem of how to effectively avoid the absorption of impurities, our study uses the wavelength, which the total lignans have higher UV absorption and the impurity has no or less absorption, as the detection wavelength of this experiment to avoid the interference of impurities. For the problem of how to purify the total lignans of Radix Isatidis more effectively, this study fully considers that the research object is the headquarters with the same basic nuclear structure, according to the principle of similarity of ultraviolet absorption of the same basic nuclear structure, so Ultraviolet spectrophotometry was used to collect the lignans components in sections, and other components of the UV absorption spectrum were discarded to purify and increase the total lignan content, which provides a basis and guarantee for the study of the pharmacological effects of the total lignans in Radix Isatidis. For the study of the pharmacological activity of the total lignans in Radix Isatidis, we first separated the total lignans from Radix isatidis and used plaque reduction experiments (CPE method is a classic method for studying viral infection experiments. This method is simple, Intuitively), the anti-RSV effects were studied from the three stages of adsorption, entry and replication of virus invading cells. The results showed that the total lignans of Radix Isatidis could significantly inhibit the formation of plaques during RSV adsorption and replication. Although this study shows that the total lignans of Radix Isatidis can exert antiviral effects during the adsorption and replication stages, it is still unknown for the mechanism of how the target exerts antiviral activity. And because this article is limited to respiratory syncytial virus, it does not mean that the total lignans are effective against all viruses. So, further research is needed.

Acknowledgment

This project was supported by National Natural Science Foundation of China (No. 81473316), Natural Science Foundation of Jiangsu Province (No.BK20151356 and BK20161320).

References

1. Yang Z, Wang Y, Zheng Z, Zhao S, Zhao J, Lin Q, et al. Antiviral activity of Isatis indigotica root-derived clemastanin B against human and avian influenza A and B viruses in vitro. *Int J Mol Med.* 2013;31(4):867-73.
2. He LW, Liu HQ, Chen YQ, Yang JY, Wang TL, Li W. Total synthesis and anti-viral activities of an extract of Radix isatidis. *Molecules.* 2014;19(12):20906-12.
3. Zhao ZJ, Wang ZW, Guo M, Du T, Shi X, Shao J, et al. Comparative study on anti-inflammatory and immunomodulation of different Isatidis Radix strains growing in Gansu Province. *Zhongguo Ying Yong Sheng Li Xue Za Zhi.* 2018;34(1):57-60.
4. Ma LN, Zhang CE, Yan D, Tan MR, Li HB, Zhang LL, et al. Screening bioactive compounds inhibiting influenza virus from isatidis radix by ultrafiltration mass spectrometry. *Zhongguo Zhong Yao Za Zhi.* 2014;39(5):812-6.
5. Lijing K, Teng W, Bradshaw JP, Zhou J, Rao P. Antiviral Decoction of Isatidis Radix (ban lan gen) Inhibited Influenza Virus Adsorption on MDCK Cells by Cytoprotective Activity. *J Tradit Complement Med.* 2012;2(1):47-51.
6. Xiao P, Huang H, Chen J, Li X. In vitro antioxidant and anti-inflammatory activities of Radix Isatidis extract and bioaccessibility of six bioactive compounds after simulated gastro-intestinal digestion. *J Ethnopharmacol.* 2014;157:55-61.
7. Zhou W, Zhang XY. Research Progress of Chinese Herbal Medicine Radix isatidis (Banlangen). *Am J Chin Med.* 2013;41(4):743-64.




Figure 3: Lignans treatment suppresses RSV F and G protein expression.




Figure 4: Inhibitory effect of lignans on RSV infection. (A) The total lignans low dose group. (B) The total lignans medium dose group. (C) The total lignans high dose group. (D) Ribavirin group.

8. Wang XL, Chen MH, Wang F, Bu PB, Lin S, Zhu CG, et al. [Chemical constituents from root of *Isatis indigotica*]. *Zhongguo Zhong Yao Za Zhi*. 2013;38(8):1172-82.
9. Cai SQ, Wang X, Shang MY. "Efficacy Theory" may help to explain characteristic advantages of traditional Chinese medicines. *China Journal of Chinese Materia Medica*. 2015;40(17):3435.
10. Zuo L, Li JB, Xu J, Yang JZ, Zhang DM, Tong YL. [Studies on chemical constituents in root of *Isatis indigotica*]. *Zhongguo Zhong Yao Za Zhi*. 2007;32(8):688-91.
11. Li L1, Qiao CZ, Li XL, Dong TY. [Study of quality control on herbal drugs and preparations of daqingye and banlangen]. *Yao Xue Xue Bao*. 1993;28(3):229-33.
12. Yang L, Wang G, Wang M, Jiang H, Chen L, Zhao F. et al. Indole alkaloid from the roots of *Isatis indigotica* and their inhibitory effects on nitric oxide production. *Fitoterapia*. 2014;95:175-81.
13. Kong W, Zhao Y, Shan L, Xiao X, Guo W. Thermochemical studies on the quantity-antibacterial effect relationship of four organic acids from *Radix Isatidis* on *Escherichia coli* growth. *Biol Pharm Bull*. 2008;31(7):1301-5.
14. Fang JG, Liu YH, Wang WQ, Xie W, Fang SX, Han HG. The anti-endotoxic effect of *o*-aminobenzoic acid from *Radix Isatidis*. *Acta Pharmacol Sin*. 2005;26(5):593-7.
15. Song QY, Zhang CJ, Li Y, Wen J, Zhao XW, Liu ZL, et al. Lignans from the fruit of *Schisandra sphenanthera*, and their inhibition of HSV-2 and adenovirus. *Phytochemistry Letters*. 2013;6(2):174-8.
16. Peng J, Fan G, Wu Y. Isolation and purification of clemastanin B and indigotcoside A from *Radix Isatidis* by high-speed counter-current chromatography. *J Chromatogr A*. 2005;1091(1-2):89-93.
17. Zhou W, Zhang X, Xie M, Chen Y, L, Y, Duan GL. Infrared-Assisted Extraction of Adenosine from *Radix Isatidis* Using Orthogonal Experimental Design and LC. *Chromatographia*. 2010;72(7-8):719-24.
18. Wu G, Wu R, Zhao W, Wang CY, Hao DD, Qu YP. Study of Water-extraction Process of Epigoinin from *Radix isatidis* by Orthogonal Experiment. *Journal of Heilongjiang Bayi Agricultural University*. 2014;6:8.
19. Luan G, Wang H, Lv H, Hu N, Suo Y, Wang X. Separation and Purification of Five Flavone Glucosides and One Lignan from *Caragana korshinskii*, Kom by the Combination of HSCCC and Semi-preparative RPLC. *Chromatographia*. 2016;79(13-14):823-31.
20. Xiong QP, Zhang QH, Xu YP, Shi YY, Zhang DY. Purification technology of epigoinin from *RADIX ISATIDIS* by macroporous resin. *Medicinal Plant*. 2011;12:49-52.
21. Cao YG, Hao Y, Li ZH, Liu ST, Wang LX. Antiviral activity of polysaccharide extract from *Laminaria japonica*, against respiratory syncytial virus. *Biomed Pharmacother*. 2016;84:1705-1710.
22. Park KI, Park HS, Kang SR, Nagappan A, Lee DH, Kim JA, et al. Korean *Scutellaria baicalensis* water extract inhibits cell cycle G1/S transition by suppressing cyclin D1 expression and matrix-metalloproteinase-2 activity in human lung cancer cells. *J Ethnopharmacol*. 2011;133(2):634-41.
23. Shi H, Ren K, Lv B, Zhang W, Zhao Y, Tan RX, et al. Baicalin from *Scutellaria baicalensis* blocks respiratory syncytial virus (RSV) infection and reduces inflammatory cell infiltration and lung injury in mice. *Sci Rep*. 2016;6:35851.