

## Clinical Image

# Gangrene of Penile Skin - A Rare Entity

Jha SK\* and Qamar A

Department of Surgery, Northern Railway Central Hospital, India

## Clinical Image

Penile gangrene is seen infrequently but is associated with significant morbidity and mortality. Gangrene of penis may be dry or infective. It is rare, fulminant but usually a localized disease of penis and scrotum with occasional extension up to abdominal wall.

A 27-year-old gentleman presented with pain, fever and brownish black discoloration of penis since 7-8 days; with history of savalon application over the penile area for hygiene purpose. There was no h/o trauma, infection or diabetes.

- General examination: febrile; no palor.
- Local examination: brownish black discoloration of penile skin extending up to penoscrotal junction without any clear line of demarcation.

Penile gangrene is rare, but grave sequential following infection or ischemia. Vascular Compromise may be caused by tourniquet syndrome; priapism; venous thrombosis and anticoagulants and can also be seen in diabetics with end stage renal disease (Figure 1). USG Doppler can provide information regarding thrombosis or rupture of cavernosal and dorsal penile vessels (Figure 2).




Figure 1: Penile gangrene.




Figure 2: USG doppler shows thrombosis or rupture of cavernosal and dorsal penile vessels.

**Citation:** Jha SK, Qamar A. Gangrene of Penile Skin - A Rare Entity. J Surg Surgic Case Rep. 2022;3(1):1023.

**Copyright:** © 2022 Sonal Kumar Jha

**Publisher Name:** Medtext Publications LLC

**Manuscript compiled:** May 24<sup>th</sup>, 2022

**\*Corresponding author:** Sonal Kumar Jha, Department of Surgery, Northern Railway Central Hospital, India, E-mail: sonaljha.jha@gmail.com