

Research Article

In Vitro Efficacy of Antimalarial Drug Primaquine against Adult Cestode and Trematode Helminthes of Sheep

Hatem A Shalaby¹, Amira H El Namaky^{1*}, Heba M Ashry¹, Faragalla El Moghazy¹, Reem OA Kamel² and Korany Farag T¹

¹Department of Parasitology and Animal Diseases, National Research Centre, Egypt

²Department of Zoology, Ain Shams University, Egypt

Abstract

The fact that the efficiency of anthelmintic drugs may diminish through ten years of use depending on various issues, one of these issues includes producers' over-dependence on such chemicals for parasites control, has dragged the concern of many researchers to use other drugs. Both scanning electron and light microscopic studies were used, for the first time, to determine the effect of antimalarial drug primaquine on some gastrointestinal helminthes of sheep following 24 h *in vitro* incubation. These results were compared with those observed on adult worms exposed to albendazole (reference drug). Two species of helminthes, *Moniezia expansa* and *Paramphistomum microbothrium*; cestoda and trematode representatives were subjected to 10 µg/ml, 20 µg/ml and 30 µg/ml of primaquine. The helminthes' tegument was affected and altered by primaquine. Furthermore, the response to primaquine action was more obvious in *P. microbothrium* than *M. expansa* adult worms. These effects involved destruction, alterations, and deformation of the tegument of *P. microbothrium*, while limited effect was observed on the tegument of *M. expansa*. This study suggested that primaquine is not only a potent antimalarial drug, but also effective trematocidal drug causing significant damage to the fluke's tegument.

Keywords: Primaquine; Antimalarial; Albendazole; *Moniezia expansa*; *Paramphistomum microbothrium*; *In vitro* effect

Introduction

Primaquine has been used since the early 1950s and is the most widespread 8-aminoquinoline antimalarial drug (WHO 2015). *In vitro*, it displayed antischistosomal activities against both juvenile and adult worms of *Schistosoma mansoni* that caused the body of parasite to be deformed in a prominent manner [1]. According to literatures primaquine was found to affect the schistosomula lysosomal acidic vesicles which responsible for endocytosis and detoxification [2]. Besides, it reduced the survival of both male and female worms and inhibited daily egg output [3]. Recently, there were no published studies concerning using primaquine as anti-trematocidal or anti-cestodal drug. On the other hand, gastrointestinal helminthes infection is the most common parasitic infection of ruminants worldwide, more particularly in third-world countries because it infects livestock to a large extent and is now well recognized as the highest disease cost to small ruminant industry [4]. Two species of helminthes have the most economic significance; the cestoda *M. expansa* and the trematode *P. microbothrium*. *M. expansa* is a very common parasite that infects

sheep, as well as goat populations and could be considered as an important problem in sheep breeding [5]. The vast majority species of paramphistomes in Egypt is *P. microbothrium* [6]. It causes economic loss greater than those caused by many other parasites [7]. As denoted by Khani et al. [8], the immature paramphistomes caused very high death, representing 80% - 90% in domesticated ruminants. The sheep industry still relies heavily on the use of anthelmintics to alleviate the infections of gastrointestinal helminthes, in spite of these drugs are expensive and usually do not block reinfection [9]. Moreover, Shalaby reported that the efficacy of anthelmintics might decrease through about ten years of use based on producers' over-dependence on such chemicals in treatment as well as poor administration practices such as under-dosing [10]. So, the need for new anthelmintics from a different chemical group in veterinary medicine is insistent. As mentioned by McKinstry that helminthes' tegument is vital for the absorptive and protective functions, the current work was undertaken to assess whether the primaquine had any effect on the tegument of both adult *M. expansa* and *P. microbothrium* following incubation *in vitro* [11]. The results were matched with those detected in the helminthes' tegument after exposure to albendazole, as it was one of the most effective of the broad-spectrum anthelmintic agents.

Materials and Methods

Drugs

Primaquine bisphosphate was obtained from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA), 10 mg/ml primaquine was prepared as a stock solution with 3ml double distilled water [2]. Albendazole (Vermizole®) was purchased from Amoun Pharmaceutical Company (El-Obour City, Cairo, Egypt).

Parasites

P. microbothrium and *M. expansa* adult worms were collected from the rumens and intestines, of naturally infected sheep slaughtered in

Citation: Shalaby HA, Namaky AHE, Ashry HM, Moghazy FE, Kamel RO, Farag TK. *In Vitro* Efficacy of Antimalarial Drug Primaquine against Adult Cestode and Trematode Helminthes of Sheep. World J Vet Sci. 2020; 2(1): 1011.

Copyright: © 2020 Hatem A Shalaby

Publisher Name: Medtext Publications LLC

Manuscript compiled: Apr 10th, 2020

***Corresponding author:** Amira H El Namaky, Department of Parasitology and Animal Diseases, National Research Centre, Giza, Egypt, Tel: 01006340534; Fax: 020 3370931; E-mail: amiraelnamaky@gmail.com

Cairo abattoir. Worms were washed in different changes of sterilized culture medium-RPMI 1640.

Anthelmintic effects of primaquine

The adult worms, after recovery, were conveyed to fresh RPMI containing 50% (v/v) heat denatured rabbit serum, 2% (v/v) rabbit red blood cells; as recommended by Ibarra and Jenkins, and primaquine at concentrations of 10 µg/ml, 20 µg/ml and 30 µg/ml [12]. Then the worms were incubated at 37°C for 24 h in an atmosphere of 5% CO₂. A reference drug group was prepared by incubating the adult worms in RPMI 1640 culture medium containing 10 µg/ml albendazole sulfoxide, (ABZ-SO) active form, for 24 h. This concentration is close to the ultimate blood levels of the sulfoxide metabolite *In vivo* [13]. A stock solution of albendazole was prepared in DMSO. Then it added to RPMI medium to give a solvent concentration of 0.1% (v/v). A normal control group was prepared by incubating worms in RPMI medium without adding drugs. Six worms from each group were examined.

Light microscopy

After incubation, 10% buffered formol saline was used to fix the adult worms. The samples were dehydrated with a graded ethanol series then embedded in paraffin. Longitudinal sections of *M. expansa* gravid segments and cross sections of *P. microbothrium* were cut with a microtome. The sections (4 µm - 6 µm thick) were stained with hematoxyline and eosin according to the method of Bancroft et al. [14]. The tegument of adult worms was studied and photographed using an Olympus CX41 microscope.

Scanning electron microscopy (SEM)

After incubation for 24 h in 30 µg/ml primaquine, the anterior end of adult *M. expansa* and intact adult *P. microbothrium* were fixed in a 3:1 mixture of 4% (w/v) glutaraldehyde in 0.12 M Millonig's buffer, pH 7.4 and 1% aqueous osmium tetroxide. After washing with distilled water, dehydration of the specimens occurred by increasing concentrations of ethanol (from 50% to 100%), and then dried in a HCP-2 critical point drying apparatus (Hitachi, Japan) using liquid carbon dioxide as a transitional medium for 15 min. The specimens were mounted on aluminum stubs and coated with gold in an ion-sputtering apparatus for 4 min. The specimens were examined using a Jeol scanning electron microscope (Jeol Corp., Mitaka, Japan) operated at 15 kV.

Results

The control untreated adult worms exhibited no loss of motility during the 24 h, the whole period of incubation. While, the adult *M. expansa* worms treated with 30 µg/ml primaquine responded less sensitive to changes in the surrounding conditions than the control worms, at a time, the adult *P. microbothrium* showed complete loss of motility. All the adult worms incubated at both 10 µg/ml and 20 µg/ml primaquine exhibited active movement throughout the incubation period of 24 h. On the other hand, those exposed to 10 µg/ml ABZ-SO (reference drug) showed paralysis or death of the adult worms. To determine the mechanisms by which primaquine, at 30 µg/ml concentration, affected the adult worm activity, the possible tissue damage induced after treatment was evaluated analyzing histological sections of the tegument of the adult worm and the structures of the tegumental surface.

Light microscopic observations

***M. expansa*:** The tegument of the control worm's gravid segments

showed features similar to that of the fresh normal specimen. Briefly, it showed an intensively stained syncytial layer; lay on a thick basement membrane of amorphous material containing granular inclusions. The basement membrane appeared to be continuous with the general filling material which lay between parenchymal cells of the interior of the proglottis. Beneath the outer tegumentary layer were a sheath of muscle fibers; an outer layer of circular muscle and an internal layer of longitudinal muscle. Those layers were followed by a sub tegumentary layer of branching parenchymal cells filled the whole space around the uterine branches (Figure 1A-C). In the primaquine treated worms, the tegument appeared to be more swollen than normal, while the underlying structures still appeared normal (Figure 1D-F). This swelling became more pronounced in ABZ-SO treated worms. In those specimens, the tegument lost its normal aspect and appeared to be extremely corrugated, swollen and pale, accompanied by the appearance of prominent wrinkles on its outer layer. In some areas,




Figure 1: Light micrographs of the tegument longitudinal section of adult *M. expansa* gravid segments.

(A-C) Normal control worms, (D-F) Following 24 hr incubation with 30 µg/ml primaquine. Note slightly swollen tegument. (G-I) Following 24 hr incubation with 10 µg/ml albendazole sulphoxide. The tegument is extremely corrugated, swollen and pale as compared with the controls. The outer tegumental layer shows prominent wrinkles (arrows) with patches of tegumental sloughing. Some areas exhibit disruption of the muscle bundles.

disruption of the muscle bundles was observed (Figure 1G-I).

***P. microbothrium*:** No significant differences in the tegumental features were noticed between normal and control worms incubated for 24 h in media free from drug. The tegument showed even and intensely stained cytoplasmic syncytial layer, which rested on basal lamina. The latter linked the tegument to the underlying two muscular layers which sent their processes outwardly to join up with the tegument. The tegumental cells located underneath the muscular layers (Figure 2A-C). The primaquine treated fluke's revealed severe tegumental disruption and sloughing of patches of the outer tegumental layer exposing the basal lamina. The muscle bundles and the parenchymal tissues showed severe degenerative changes (Figure 2D-F). Degenerative changes of the outer tegumental layer were also apparent following treatment with ABZ-SO, but the muscles underlying the tegument still exhibited a normal appearance (Figure 2G-I).

Scanning electron microscopic observations

***M. expansa*:** A globular scolex provided with four oval suckers radially located at its peripheral margin was observed at the anterior end of the control adult *M. expansa*. The tegument behind the suckers

had no in folding (Figure 3A). The scolex appeared to be more swollen than the control with narrowing of the sucker's opening following treatment with primaquine. The tegument lost its normal aspect showing corrugated tegumental surface (Figure 3B). Following ABZ-SO treatment, the adult cestode exhibited swollen scolex with severely folded tegument around the suckers so that the contractions of their openings were obvious (Figure 3C).




Figure 2: Light micrographs of the tegument cross section of adult *P. microbothrium*. (A-C) Normal control fluke. (D-F) Following 24 h incubation with 30 µg/ml primaquine. Note severe tegumental disruption and sloughing of patches of the outer tegumental layer exposing the basal lamina. The muscle bundles and the parenchymal tissues show severe degenerative changes. (G-I) following 24 h incubation with 10 µg/ml albendazole sulphoxide. Note degenerative changes of the outer tegumental layer. T: Tegument; BL: Basal Lamina; M: Muscular layer; TC: Tegumental Cell; P: Parenchyma




Figure 3: Scanning electron micrographs of the anterior end of adult *M. expansa*. (A) Normal control worm showing a globular scolex provided with four oval suckers and smooth tegument. (B) Following 24 h incubation with 30 µg/ml primaquine. The scolex becomes more swollen than the control with narrowing of the sucker's opening and corrugation of the tegumental surface. (C) Following 24 h incubation with 10 µg/ml albendazole sulphoxide. The scolex is swollen with severely folded tegument around the suckers and contraction of their openings.

P. microbothrium: In the control, the adult fluke had a pear-shaped body with a narrow anterior end and broader posterior end (Figure 4A,B). The oral sucker located near the anterior tip, was transversely elongated oval, whereas the acetabulum was positioned close to the posterior tip. The tegument covering the body was composed of transverse folds alternating with grooves (Figure 4C). Aggregation of dome-shaped papillae arranged in rows was clearly visible in these anterior folds. In almost all specimens examined of the primaquine treated flukes, distortion of both oral sucker and acetabulum was

extreme to such extent that little distinguishable structure remained (Figure 4D,E). Apparently, patches of thickened tegument were seen throughout the posterior region around the acetabulum. At high magnification, the tegumental thickening was observed due to a layer of syncytial secretion overlying the surface of the tegument. Also, this syncytial secretion was extended to the mid-body region of the fluke so that the tegumental transverse folds could not be recognized (Figure 4F). At high magnification, neat and round holes were noticeable at the surface (Figure 4F). In ABZ-SO treated flukes, the tegumental alterations appeared similar to that of the primaquine treated flukes. Yet, a number of pits caused by rupture of papillae were observed at higher magnification of the tegumental surface at the anterior end of the fluke (Figure 4G). Acetabulum showed considerable damage; the texture of the tegument was distorted and the papillae were lost (Figure 4H). The tegumental surface at the mid-body region exhibited severe blebbing (Figure 4).

In all experiments of scanning electron microscopic studies, no significant differences in the tegumental features were seen between control and normal fresh worms incubated for 24 hr in media free from drug.




Figure 4: Scanning Electron Micrographs (SEMs) of the adult *P. microbothrium*. (A-C) Normal control fluke. (A) SEM of the anterior end showing transversely elongated oral sucker. (B) SEM of the posterior end showing the acetabulum. (C) SEM of the mid-body region of the fluke's body exhibiting smooth transverse folds alternating with grooves. (D-F) Following 24 hr incubation with 30 µg/ml primaquine. (D,E) SEMs of the anterior and posterior ends reveal distortion of both oral sucker and acetabulum as well as a layer of syncytial secretion overlying the surface of the tegument throughout the posterior region around the acetabulum. (F) SEM of the mid-body region showing neat rounded holes. (G-I) Following 24 hr incubation with 10 µg/ml albendazole sulphoxide. (G) SEM of the anterior end showing a number of pits caused by rupture of papillae. (H) SEM of the acetabulum exhibiting distortion of the tegument texture. (I) SEM of the mid-body region exhibiting severe blebbing.

Discussion

The current study demonstrated the comparative morphological effects of primaquine and ABZ-SO (reference drug) against *M. expansa* and *P. microbothrium* adult worms; cestode and trematode representatives. This is the first study demonstrating the *in vitro* effects of primaquine on some gastrointestinal helminthes of sheep.

Remarkably, primaquine is the fourth antimalarial drug showing anthelmintic properties after artemisinin (artemether and artesunate), trioxolanes and mefloquine [15-18]. In this study, the response to primaquine action was more potent in *P. microbothrium* than *M. expansa* adult worms. During the whole period of the experiments, the 30 µg/ml primaquine treated *M. expansa* showed a slower rate of activity than the controls and none of the treated cestodes died. While, 30 µg/ml primaquine treated *P. microbothrium* and ABZ-SO treated worms showed complete loss of motility. These observations might refer to the superiority of primaquine in killing the trematodes that might be appropriate *in vivo* to drive the worms out from the host's gastrointestinal tract, as had been illustrated for albendazole [19]. Indeed, in the present study, the data of the tegument histological observations of *P. microbothrium* after *in vitro* administration of primaquine were similar to that induced by ABZ-SO, and more severe than that were observed in the primaquine treated *M. expansa*. However, the assessment of drug-derived effects was essentially depended upon electron micrographs, rather than light micrographs. Since, electron micrographs elucidated the detailed morphology and different changes of the worm's tegument permitting the interpretation of its functionality. In this aspect, the tegumental distortion of *P. microbothrium* including the oral sucker and the acetabulum, as well as the tegumental thickening throughout the mid-body region of the fluke; that was marked with neat round holes, were observed during the *in vitro* action of primaquine. The surface changes were more severe than those observed in the primaquine treated *M. expansa*, in which limited tegumental swelling had occurred in their scolices. The tegumental distortion has been seen in the adult flukes treated with ABZ-SO. Previous studies had shown that the albendazole possessed a broad spectrum activity against all classes of parasitic helminthes. This drug had been recorded to induce tegumental disruption and muscular degeneration by binding specifically to β -tubulins, thereby inhibiting polymerization and functioning of the cellular motor proteins [20,21]. In the current study, ABZ-SO showed a potential *in vitro* effect against *P. microbothrium* and *M. expansa*, where the tegument of the adult worms was severely distorted and tegumental blebs appeared especially at the mid-body region of the flukes. Similar findings were reported for, biologically related trematodes, *Cotylphoron cotylphorum* [22] and *Fasciola hepatica* treated *in vitro* with 10 µg/ml of albendazole for 12 hours [22,23]. The thickening of the tegument without bleb formation was observed in the primaquine treated flukes, but the surface was marked with neat, round holes that might be resulted from rupture of blebs. These tegumental alterations were also reported for *P. microbothrium* and *F. gigantica* treated *in vitro* with, an antimalarial drug, artemether [17,24]. The tegumental alterations; including swelling, blebbing that was later disrupted causing erosion, were common features of drug-treated trematodes and cestodes after exposure to anthelmintics [25]. Additionally, the tegumental thickening was probable to represent part of a stress response on the part of the fluke and had been observed in another anthelmintic studies on *F. hepatica* during the early stages of drug action [26-28]. In an effort to maintain the safety of the apical membrane, the layer was formed by the accelerated release of tegumental secretions at the surface.

Our observations on primaquine treated worms pointing to the tegument as the main interface for drug uptake. Much of the literature suggested trans-tegumental uptake might play a more significant role in drug entrance into trematode and cestode parasites [29]. However, in this study, the *P. microbothrium* tegument was severely affected

than that of *M. expansa* following their exposure to primaquine. Traditionally, the complex trematode and cestode tegument was believed to act as absorptive surface [30]. *P. microbothrium* exhibited a high degree of corrugation comprising of alternating grooves and folds increasing the surface area of absorption by the tegument. The higher absorptive ability in the tegument of *P. microbothrium* reflected on the higher level of disruption to the flukes when exposed to primaquine. A recent published study showed that 24 hr after incubation in primaquine at the concentration of 20 µg/ml resulted in extensive damage of, a biologically related trematode, *Schistosoma mansoni* including degeneration of both the tegumental and subtegumental layers [1]. Also, Holtfreter et al. [2] suggested a strong impact of the primaquine on the tegument of *S. mansoni*.

Conclusion

The present study has suggested that primaquine is not only potent antimalarial drug, but also effective trematocidal drug causing significant damage of the fluke's tegument. It means that the parasite's first and main line of defense is destroyed allowing the drug potential access to other, internal tissues of the fluke, which will lead to more widespread damage. However, several questions remain to be answered, for example the mechanism by which primaquine exerts its effect on the parasite's tegument, and why the intensity of primaquine-induced damage in *P. microbothrium* is greater than that in *M. expansa*.

References

- Kamel ROA, Bayaomy FEA. Ultrastructural alterations in *Schistosoma mansoni* juvenile and adult male worms after *in vitro* incubation with primaquine. Mem Inst Oswaldo Cruz. 2017;112:247-54.
- Holtfreter MC, Loebermann M, Klammt S, Sombetzki M, Bodammer P, Riebold D, et al. *Schistosoma mansoni*: schistosomicidal effect of mefloquine and primaquine *in vitro*. Exp Parasitol. 2011;127(1):270-6.
- Mitsui Y, Aoki Y. *In vitro* effects of current antimalarial drugs on the survival of paired *Schistosoma mansoni* adult worms and their egg production. Trop Med Health. 2010;38(2):69-73.
- Bedada H, Gizaw F, Negash W. Preliminary study on small ruminant GIT helminthiasis in select arid and semi-arid pastoral and agro-pastoral areas of Afar region, Ethiopia. Am J Parasitol. 2018;1:1-9.
- Maziad SA, El-Nemr HI. The endoparasites of sheep and goats, shepherd in North Sinai Governorate, Egypt. J Egypt Soc Parasitol. 2002;32:119-26.
- Hiekal F, Hilali M. Scanning electron microscopy of the tegument of *Paramphistomum microbothrium* Fiscoeder, 1901 and *Cotylphoron cotylphorum* (Fiscoeder 1901) in Egypt. Arab Gulf J Sci Res. 1993;11:105-13.
- Hanna REB, Williamson DS, Mattison RG, Nizami WA. Seasonal reproduction in *Paramphistomum epiclitum* and *Gastrothylax crumenifer*, rumen paramphistomes of the Indian water buffalo, and comparison with the biliary paramphistome *Gigantocotyle explanatum*. Int J Parasitol. 1988;18:513-21.
- Khani UJ, Tanveerl A, Maqbool A, Masood S. Epidemiological studies of paramphistomosis in cattle. Vet Arc. 2008;78(3):243-51.
- Fikru R, Teshale S, Reta D, Yosef K. Epidemiology of gastrointestinal parasites of ruminants in western Oromia, Ethiopia. Int J Appl Res Vet M. 2006;4:51-7.
- Shalaby HA. Anthelmintics resistance; How to overcome it? Iran J Parasitol. 2013;8(1):18-32.
- McKinstry B, Brennan GP, Halferty L, Forbes AB, Fairweather I. Ultrastructural changes induced in the tegument and gut of *Fasciola hepatica* after *in vivo* and *in vitro* drug treatment with nitroxylnil (Trodox). Parasitol Res. 2007;101(4):929-41.
- Ibarra OF, Jenkins DC. An *in vitro* screen for new fasciolicidal agents. Z Parasitenkd. 1984;70:655-61.

13. Fetterer R, Rew RS, Knight R. Comparative efficacy of albendazole against *Fasciola hepatica* in sheep and calves: relationship to serum drug metabolite levels. *Vet Parasitol.* 1982;11(4):309-16.
14. Bancroft JD, Stevens A, Turner DR. *Theory and practice of histological techniques*, 4th ed. Churchill Livingstone, New York. 1996.
15. Keiser J, Morson G. *Fasciola hepatica*: surface tegumental responses to *in vitro* and *in vivo* treatment with the experimental fasciolicide OZ 78. *Exp Parasitol.* 2008;119(1):87-93.
16. Keiser J, Shu-Hua X, Tanner M, Utzinger J. Artesunate and artemether are effective fasciolicides in the rat model and *in vitro*. *J Antimicrob Chemother.* 2006;57(6):1139-45.
17. Shalaby HA, El Namaky AH, Kamel ROA. *In vitro* effect of artemether and triclabendazole on adult *Fasciola gigantica*. *Vet Parasitol.* 2009;160(1-2):76-82.
18. Shalaby HA, El Namaky AH, Kamel ROA. *In vitro* tegumental alterations on adult *Fasciola gigantica* caused by mefloquine. *J Parasit Dis.* 2016;40(1):145-51.
19. Tinar R, Coskun S Z, Dogan H. Efficacy of albendazole against nematode and trematode infection in sheep. *Veteriner Fakultesi Dergisi Uludag Univ.* 1988;7:117-23.
20. Lacey E. The role of the cytoskeletal protein tubulin in the mode of action and mechanism of drug resistance to benzimidazoles. *Int J Parasitol.* 1988;18(7):885-936.
21. Ahmed M, Nizami WA. *In vitro* effect of mebendazole on the carbohydrate metabolism of *Gigantocotyle explanatum* (Trematoda:Digenea). *Indian J Parasitol.* 1991;15:19-26.
22. Radwan NA, Khalil AI, Wahdan AE. *In vitro* evaluation of antihelminthic activity of *Allium sativum* against adult *Cotylophoron cotylophorum* (Paramphistomidae). *PUJ.* 2012;5:135-46.
23. Buchanan JE, Fairweather I, Brennan GP, Trudgett A, Hoey EM. *Fasciola hepatica*: Surface and internal tegumental changes induced by treatment *in vitro* with the sulphoxide metabolite of albendazole (Valbazen). *Parasitol.* 2003;126:141-53.
24. Shalaby HA, El Namaky AH, Kamel ROA, Derbala AA. Tegumental surface changes in adult *Paramphistomum microbothrium* (Fischöeder 1901) following *in vitro* administration of artemether. *J Helminthol.* 2010;85(2):115-22.
25. Meaney M, Fairweather I, Brennan GP, Ramasamy P, Subramanian PB. *Fasciola gigantica*: tegumental surface alterations following treatment *in vitro* with the sulphoxide metabolite of triclabendazole. *Parasitol Res.* 2002;88(4):315-25.
26. Stitt AW, Fairweather I. *Fasciola hepatica*: tegumental surface changes in adult and juvenile flukes following treatment *in vitro* with the sulphoxide metabolite of triclabendazole (Fasinex). *Parasitol Res.* 1993;79(7):529-36.
27. Anderson HR, Fairweather I. *Fasciola hepatica*: ultrastructural changes to the tegument of juvenile flukes following incubation *in vitro* with deacetylated (amine) metabolite of diamphenethide. *Int J Parasitol.* 1995;25(3):319-33.
28. McKinstry B, Brennan GP, Halferty L, Forbes AB, Fairweather I. Ultrastructural changes induced in the tegument and gut of *Fasciola hepatica* after *in vivo* and *in vitro* drug treatment with nitroxylnil (Trodax). *Parasitol Res.* 2007;101(4):929-41.
29. Toner E, McConvery F, Brennan GP, Meaney M, Fairweather I. A scanning electron microscope study on the route of entry of triclabendazole into the liver fluke, *Fasciola hepatica*. *Parasitol.* 2009;136(5):523-35.
30. Bashtar A, Hassanein M, Abdel-Ghaffar F, Al-Rasheid KAS, Hassan S, Mehlhorn H, et al. Studies on moniezia of sheep I. Prevalence and antihelminthic effects of some plant extracts, a light and electron microscopic study. *Parasitol Res.* 2011;108:177-86.