

Research Article

Serological Evaluation of Toxocarosis in Amedyia District, Duhok Governorate, Kurdistan Region, Iraq

Haifa G Ismail¹ and Adel T Al-Saeed^{2*}

¹Department of Microbiology, Amedyia Technical Institute, Polytechnic University, Iraq

²Department of Medical Microbiology, College of Medicine, University of Duhok, Iraq

Abstract

The ova of *Toxocara canis* is common environmental contaminants of human habitation, due to the fact that dogs serve as final hosts. The presence of *Toxocara* ova in the soil which considerable as a risky factor for human public health. Humans, particularly children, frequently ingest these ova accidentally and infected with disease. Infection in humans, in contrast to their definitive hosts, remains unusual host, often resulting in disease caused by the migrating larval stages without development to adult stage. This study is the first study in Duhok Governorate and Kurdistan region to determine the seroprevalence of toxocarosis among human population and the relation of the associated factors. A total of 600 blood samples were collected from children and adults of different ages (5-70) years and both genders. Blood samples were collected in a gel tubes, then centrifuged for isolation of sera. The sera were kept at -20°C until used for detection of anti *Toxocara canis* IgG antibodies using ELISA. Out of 600 serum samples 38 (6.3%) were seropositive. The rate of infection was higher in females 22/285 (7.7%) than males 16/315 (5.1%). The individuals belong the age group less than 5 years were more prevalent (21.4%) than other groups followed by age group (5-14) year with infection rate (9.6%). Toxocarosis is prevalent in adults and children of Amedyia district. These results require periodical studies on the rate of infection and associated risk factors in other areas of Duhok Governorate and Kurdistan region.

Keywords: *Toxocara canis*; ELISA; Children; Larva migrans; Duhok; Kurdistan

Introduction

Toxocara species are roundworms from Ascaridae family, such as *Toxocara canis*, *Toxocara cati*, *Toxocara malaysiensis*, and *Toxascaris leonina*, are the most common zoonotic gastrointestinal helminths [1-4], that habit the intestine of dogs and cats throughout the world [5,6].

The ova of *Toxocara* species are common environmental contaminants of human habitation, due to the fact that the dogs and cats serve as final hosts and the presence of *Toxocara* ova in the soil which considerable as a risky factor for public health [7]. These ova may remain viable in the soil for months or even years [8]. Children's play habits and their outdoor activities put them at higher risk for infection [9,10].

Humans, particularly children, frequently ingest these ova accidentally and become infected. Infection in humans, in contrast to their definitive hosts, often resulting in disease caused by the migrating larval stages. Visceral and ocular larva migrans are two clinical manifestations that result due to migration of larvae [11,12].

The Enzyme-Linked Immunosorbent Assay (ELISA), which

employs antigens secreted by the second-stage larvae of *T. canis*, has sufficient specificity to be the reliable indirect test for diagnosing this infection [13,14].

Although comprehensive studies on *Toxocara canis* were conducted in the different regions of the world, little studies were conducted in Iraq. The aim of current study was to determine the seropositivity of *Toxocara canis* in children and adults and relation to the associated factors with infection in Amedyia District, Duhok Governorate, Kurdistan Region.

Materials and Methods

A total of 600 blood samples were collected from children and adults of different ages (5-70) years and both sexes.

The present study conducted in different areas of Amedyia district in Duhok Governorate to determine the seropositivity rate of toxocarosis using ELISA as qualitative screening and determine the relation of the risk factors associated to the infection such as contact with dogs and soil, food habitation, residency, life style, and socioeconomic status. Blood samples were collected in gel tubes, then centrifuged for isolation of sera. The sera were kept at -20°C until used for detection anti *Toxocara canis* IgG antibodies using a commercial ELISA Kit (DEMDITEC Diagnostics GmbH EN ISO 9001 certified company) according to instructions of the company. Analysis of data was carried out using the Chi Square test and Statistical Package for Social Science (SPSS).

The protocol of the current study was approved by the researches Ethics Committee of the College of Medicine, Duhok University and Duhok Directorate of Health.

Results

A total of 600 subjects examined in the present study, 357 (59.5%) were children and 243 (40.5%) were adults. Regarding to gender, 285 (47.5%) were female and 315 (52.5 %) were male. A total of 336 (56.0%)

Citation: Ismail HG, Al-Saeed AT. Serological Evaluation of Toxocarosis in Amedyia District, Duhok Governorate, Kurdistan Region, Iraq. Med Life Clin. 2022; 4(1): 1038.

Copyright: © 2022 Haifa G Ismail

Publisher Name: Medtext Publications LLC

Manuscript compiled: Aug 26th, 2022

***Corresponding author:** Adel T Al-Saeed, Department of Medical Microbiology, College of Medicine, University of Duhok, Duhok Governorate, Kurdistan Region, Iraq, E-mail: adelalsaeed@uod.ac; qaisibraheem@uod.ac

(%) were lived in rural area, 182 (30.3 %) in urban area and 82 (13.7%) in suburban area. Overall, 536 (89.3) were contact with dogs, and 64 (12.5%) were not contact. The other socio-demographic characters and associated factors including age groups were listed in (Tables 1-3).

Table 1: Frequency distribution of sociodemographic characteristics and associated factors of study participants (no. 600).

Variable		Frequency	Percent
Gender	Male	315	52.5
	Female	285	47.5
Age Group (Year)	< 5	14	2.3
	5-14	343	57.2
	15-24	46	7.7
	25-44	154	25.7
	45 and above	43	7.2
Residency	Rural	336	56
	Suburban	82	13.7
	Urban	182	30.3
Contact with dogs	No	64	12.5
	Yes	536	89.3
Socio-economic status	Poor	152	25.3
	Moderate	86	14.3
	Good	362	60.3
Total		600	100

Table 2: Seroprevalence of toxocarosis among population according to age and sex in Amedyia district, Duhok governorate.


Variables	No. of Examined	No. (%) +ve	No. (%) -ve	df	P-value
Age group (Year)					
< 5	14	3 (21.4)	11 (78.6)	4	0
5 - 12	343	33 (9.6)	310 (90.4)		
15 - 24	46	0	46 (100)		
25 - 44	154	1 (0.6)	153 (99.4)		
45 and above	43	1 (2.3)	42 (94.7)		
Sex					
Gender	Male	16(5.1)	299 (94.9)	1	0.185
	Female	22 (7.7)	263 (92.3)		

Table 3: Seroprevalence of toxocarosis associated factors in Amedyia district.

Variables	No. of Examined	No. (%) +ve	No. (%) -ve	df	P-value
Occupation					
House wife	80	1 (1.3)	79 (98.7)	3	0
Student	342	26 (7.6)	316 (82.4)		
Child	44	10 (22.7)	34 (77.3)		
Unemployed	134	1 (0.7)	153 (99.4)		
Education					
Illiterate	171	2 (1.2)	79 (98.7)	3	0.001
Primary school	190	22 (11.6)	316 (82.4)		
Secondary school	221	14 (6.3)	34 (77.3)		
High education	18	0 (0)	153 (99.4)		
Socioeconomic					
Poor	152	4 (2.6)	148 (97.4)	2	0.045
Moderate	362	25 (6.9)	337 (93.1)		
Good	86	9 (10.5)	34 (89.5)		
Residency					
Rural	336	38 (11.3)	298 (88.7)	2	0
Suburban	82	0 (0)	82 (100)		
Urban	182	0 (0)	182 (100)		
Contact with dogs					
No	63	0 (0)	63 (100)	1	0.029
Yes	536	38 (7.1)	498 (92.9)		

The result has been shown that the total rate of infection among examined population (38, 6.3%) were seropositive to anti-*Toxocara* antibodies using ELISA, while 562 (93.7%) were seronegative (Figure 1).

It is obvious from (Table 2) that the individuals belong the age group less than 5 years were more prevalent 3/14 (21.4%) than other



age groups followed by age group ranging from (5-14) year-old with infection rate 33/343 (9.6%). In addition to that there was no any case reported in age group ranging from (15-24) year old. Statistical analysis revealed significant difference between the infection and different age groups.

The rate of infection was higher in females 22/285 (7.7%) than males 16/315 (5.1%) but the statistical analysis revealed no significant difference between the infection and gender.

Regarding to the occupation of examined population, the result shows that the children were more prevalent 10/44 (22.7%) followed by students 26/342 (7.6 %). Statistical analysis revealed significant difference between occupation and infection (Table 3).

It is obvious from the results that highest rate of infection 22/190 (11.6%) was among subjects that their parent whom educational status was primary school, followed by 14/221 (6.3%) and 2/171 (1.2%) among parent whom educational status was secondary school and illiterate respectively, while there was no any case recorded among parent whom educational status was higher education. Statistical analysis of the results showed the presence of highly significant difference between all educational statuses.

The present study found that more prevalent rate among group with good socioeconomic status 9/86 (10.5%) when compared with moderate 25/362 (6.9%) and poor socioeconomic status 4/152 (2.6%). Statistical analysis revealed significant difference.

All infected cases were recorded from rural areas 38/336 (11.3%) in relation to residency of examined individuals and statistical analysis revealed high significant difference.

Discussion

The present work reports for the first time serological proven human toxocarosis among population in Amedyia district, Duhok Governorate, Kurdistan region of Iraq. Toxocarosis is common in developing countries and some studies found that high frequency of human infection by *Toxocara* larvae [15-17].

Different prevalence rates of *Toxocara* infection were recorded in different countries ranging from 3% to 86% [18]. In the current study the results show agreement with this range of 10 infection when found that the total rate of toxocarosis among examined population was 6.3% using ELISA. The results of the present study were dissimilar to that have been observed by Al-Saeed et al. [17] in Mosul city (Iraq), when reported high seropositive rate 30.8% in examined children. Unexpected results were found in Fars Province (Iran) related to

toxocarosis [18], they recorded high rate of infection (30.1%) of urban children in comparison to rural children (20.2%).

A study conducted by Roldán et al. [19] revealed that low seropositive rate 1.39% in examined children of the Amazonian city (Brazil), this result was in contrast with the result of the present study when reported that the seropositivity in children aged <5 years was 21.9%. In western Iran, they found that high rate 25% of school children between (7-14) years ages were seropositive [20].

A study conducted by Jalali et al. [21] in Shiraz, southern Iran, reported (6.4%) seropositive rate in rural area among individuals with no clinical signs and (23.3%) in individuals with clinical signs but there was no significant correlation was reported between contact with dogs and *Toxocara* infection. On the other hand, high prevalence rate 76.6% of toxocarosis among (7-12) year old children in Taiwan, the reason was a significant correlation between histories of raising dogs and seropositivity [22]. While in Nigeria reported 30% seropositivity among individuals aged (2-21) year but showed that contact with dogs was not an important factor for *Toxocara* infection [23]. A study conducted by Pierangeli NB et al. [24] reported that 4.8% of healthy blood donors and 1.2% of children were infected with toxocarosis in Argentina.

Researchers showed that low rate of infection 2% in Indian children by serological examination [25]. The rate of seropositivity found among children in the present study 38/600 (6.3%) is consistent with reports conducted in children's groups (26.6%) from Egypt [26], Spain 27.2% [27], Nigeria 29.8% [28], Turkey 25.9% [29], Bolivia 24.8% [30].

The differences in seropositive rates may be due to variation in the climatic conditions such as temperature and humidity which effect on the survival and viability of *Toxocara* ova in the environment, life and eating habits (eating unwashed raw vegetables), occupation and personal hygiene.

The higher seroprevalence of anti *Toxocara* IgG antibodies in older age groups was due to their longer exposure to risk factors associated to the infection, while unexpected results in the current study were found in old age group that lowest infection rate with toxocarosis, the reasons might be due to the life style, social behaviour, socioeconomic status number, spreading and contact with dogs.

The present study reported that infection rates in females were slightly higher than males. Dissimilar results were found in other studies that males with highest infection rate compared with females [20,31].

The authors suggested that difference might be due to the fact that outdoor activities were restricted in girls and most societies due to some social and religious restrictions, females are relatively stay indoors while males have to work in outdoor.

In conclusion, toxocarosis is endemic in Amedyia district of Duhok Governorate, Kurdistan region of Iraq. These results require periodical studies on the rate of infection and associated risk factors in other areas of Duhok Governorate and Kurdistan Region.

References

- Filho FA, Chieffi PP, Correa CRS, Camargo ED, da Silveira EPR, Aranha JJB, et al. Human toxocarosis: a seroepidemiological survey in the municipality of campinas (SP), Brazil. *Rev Inst Med Trop Sao Paulo.* 2002;44(6):303-7.
- Bass JL, Mehta KA, Glickman LT, Eppes BM. Clinically inapparent toxocara infection in children. *N Engl J Med.* 1983;308(12):723-4.
- Beaver PC. Larva migrans. *Exp Parasitol.* 1956;5(6):587-621.
- Buijs J, Borsboom G, Renting M, Hilgersom WJ, van Wieringen JC, Jansen G, et al. Relationship between allergic manifestations and toxocara seropositivity: a crosssectional study among elementary school children. *Eur Respir J.* 1997;10(7):1467-75.
- Chorozy ML, Richardson DJ. A survey of environmental contamination with ascarid ova, wallingford, connecticut. *Vector Borne Zoonotic Dis.* 2005;5(1):33-9.
- De Andrade Lima Coelho R, Carvalho LBJr, Perez EP, Araki K, Takeuchi T, Ito A, et al. Prevalence of toxocarosis in northeastern Brazil based on serology using recombinant toxocara canis antigen. *Am J Trop Med Hyg.* 2005;72(1):103-7.
- De Savigny DH, Voller A, Woodruff AW. Toxocarosis: serological diagnosis by enzyme immunoassay. *J Clin Pathol.* 1979;32(3):284-8.
- Despommier D. Toxocarosis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clin Microbiol Rev.* 2003;16(2):265-72.
- Deutz A, Fuchs K, Auer H, Kerbl U, Aspöck H, Köfer J. Toxocara-infestations in Austria: a study on the risk of infection of farmers, slaughterhouse staff, hunters and veterinarians. *Parasitol Res.* 2005;97(5):390-4.
- Figueiredo SDP, Taddei JAAC, Menezes JJC, Novo NE, Silva EOM, Cristoao HLG, et al. Estudo clinic-epidemiologico da toxocarose em populacao infantil. *J Pediatr.* 2005;81(2):126-32.
- Gamboa MI. Effects of temperature and humidity on the development of eggs of toxocara canis under laboratory conditions. *J Helminthol.* 2005;79(4):327-31.
- Glickman LT, Schantz PM. Epidemiology and pathogenesis of zoonotic toxocarosis. *Epidemiol Rev.* 1981;3(1):230-50.
- Glickman LT, Grieve RB, Lauria SS, Jones DL. Serodiagnosis of ocular toxocarosis: a comparison of two antigens. *J Clin Pathol.* 1985;38(1):103-7.
- Habluetzel A, Traldi G, Ruggieri S, Attili AR, Scuppa P, Marchetti R, et al. An estimation of toxocara canis prevalence in dogs, environmental egg contamination and risk of human infection in the Marche regions of Italy. *Vet Parasitol.* 2003;113(3-4):243-52.
- Jacob CMA, Pastorino AC, Peres BA, Melo EO, Okay Y, Oselka G. Clinical and laboratorial features of visceral toxocarosis in infancy. *Rev Inst Med Trop Sao Paulo.* 1994;36(1):19-26.
- Alonso JM, Steun M, Chamorro MC, Bojanich MV. Contamination of soils with eggs of toxocara in a subtropical city in Argentina. *J Helminthol.* 2001;75(2):165-8.
- AL-Saeed WM, Al-Dabbagh NY, Mahmood HJ. Serological study of toxocarosis in children in Mosul province. *Med J Babylon.* 2009;6(3-4):455-64.
- Abdi J, Darabi M, Sayehmiri K. Epidemiological situation of toxocarosis in Iran: meta-analysis and systematic Review. *Pak J Biol Sci.* 2012;15(22):1052-5.
- Roldán WH, Cavero YA, Espinoza YA, Jiménez S, Gutiérrez CA. Human toxocarosis: a seroepidemiological survey in the Amazonian city of Yurimaguas, Peru. *Rev Inst Med Trop Sao Paulo.* 2010;52(1):37-42.
- Fallah M, Azimi A, Taherkhani H. Seroprevalence of toxocarosis in children aged 1-9 years in western Islamic Republic of Iran, 2003. *East Mediterr Health J.* 2007;13(5):1073-7.
- Jalali A, Mehrabani D, Sadjadi SM, Orryan A, Khosravi M. Visceral larva migrans and ocular larva migrans as dangerous disease in Shiraz, Southern Iran. *The Middle East J Emerg Med.* 2004;4:115-6.
- Fan CK, Lan HS, Hung CC, Chung WC, Liao CW, Du WY, et al. Seroprevalence of toxocara canis infection among mountain aboriginal adults in Taiwan. *Am J Trop Med Hyg.* 2004;71(2):216-21.
- Ajanusi OJ, Asiribo OE. A ten-year analysis of canine hookworm and toxocara canis infections in Zaria, Nigeria. *J Trop Biosci.* 2004;4:23-6.
- Pierangeli NB, Giayetto AL, Manacorda AM, Barbieri LM, Soriano SV, Veronesi A, et al. [Estacionalidad de parásitos intestinales en suelos periurbanos de la ciudad de

- Neuquén, Patagonia, Argentina]. *Trop Med Int Health*. 2003;8(3):259-63.
25. Malla N, Aggarwal AK, Mahajan RC. A serological study of human toxocariasis in north India. *Natl Med J India*. 2002;15(3):145-7.
26. Oteifa NM, Moustafa MA, Elgozamy BM. Toxocariasis as a possible cause of allergic disease in children. *J Egypt Soc Parasitol*. 1998;28(2):365-72.
27. Ruiz de Ybanez MR, Garijo MM, Alonso FD. Prevalence and viability of eggs of toxocara spp. and toxocaris leonine in public parks in eastern Spain. *J Helminthol*. 2001;75(2):169-73.
28. Amodi NO. Prevalence of parasite eggs and cyst on the Nigerian currency (Naira) in Zaria. Unpublished MSc thesis. Department of Biological Sciences, Ahmadu Bello University, Zaria. 2000;Pp 21.
29. Kaplan M, Kalkan A, Hosoglu S, Kuk S, Özden M, Demirdag K, et al. The frequency of toxocara infection in mental retarded children. *Mem Inst Oswaldo Cruz*. 2004;99(2):121-5.
30. Nicoletti A, Bartoloni A, Reggio A, Bartalesi F, Roselli M, Sofia V, et al. Epilepsy, cysticercosis, and toxocariasis: a population-based case-control study in rural Bolivia. *Neurology*. 2002;58(8):1256-61.
31. Matos CMF, Militao DNA, Brum MAR, Tundisi RN. Presence of anti-toxocara antibodies in children selected at hospital universitario, campio grande, Brazil. *Rev Inst Med Trop S Paulo*. 1997;39(1):45-51.