

## Research Article

# Toxicity of Mareb Crude Oil on Intertidal Clam *Tivela Ponderosa* and its Effect on Oxygen Consumption under Laboratory Conditions

Aziz S. Dobian<sup>1</sup> and Hagib A. Al-Hagibi<sup>2\*</sup>

Department of Earth and Environmental Sciences, Sana'a University, Yemen

## Abstract

The impact of chronic exposure of Mareb Crude Oil on the rate of oxygen consumption of the clam, *Tivela Ponderosa* was studied in the laboratory. The bivalve *T. ponderosa* is exposed to (0.5, 1 and 1.5) ml/l of Mareb Crude Oil. The oxygen consumption of bivalves *T. ponderosa* was controlled hourly at 5<sup>th</sup> and 10<sup>th</sup> day (every two and six hours). After 5 and 10 days of exposure to crude oil concentrations, the average oxygen consumption in the clam after 5 days of exposure was (0.131, 0.135, 0.141, 0.319 ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup> every two hours) and (0.121, 0.124, 0.137, 0.247 ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup> every six hours) and after 10 days it was (0.222, 0.561, 0.946, 1.117 ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup> every two hours) and (0.126, 0.432, 0.573, 0.632 ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup> every six hours). It was observed that the rate of oxygen consumption fluctuated with an increase in the exposure period. The increase in oxygen consumption in the treated clams can be explained by the high metabolic activity of the organisms due to the stress imposed by the pollutant. On the other hand, when increasing the exposure period for pollution substance decrease in oxygen consumption in the clams exposed to toxic substance concentrations. The possible reason for the latter is the insensitivity of the clams especially in high concentration of pollution substances.

**Keywords:** Toxicity; Oxygen consumption; Condition index; Crude oil; Clam *Tivela ponderosa*

## Introduction

The rate of oxygen consumption varies with changes in the environmental and physiological conditions. Changes in physiological activities of organisms serve as an indicator of sub lethal effects of pollutants on organisms [1,2]. Thus, it is an indicator to determine the degree of stress caused by changes in the environment due to various natural and man-made perturbations.

Deshmukh [3] studies the changes in oxygen consumption by the clam, *Meretrix* exposed to various changes in natural conditions (temperature, salinity, etc.). Rate of oxygen consumption has been used as available tool by many workers to assess stress; since it is an index of energy expenditure to meet the demands of environmental alterations [4-6] Most of the vital activities in bivalves are regulated by Neuro-endocrine centers. The respiratory rate data of the animals reflect their general metabolic rate. The aim of this study by using bivalves (*T. ponderosa*) as bio indicator to determine the effect of toxic for Mareb Crude Oil on oxygen consumption rate in bivalve mollusks. In addition to study of the condition index of the exposed bivalve.

## Materials and Methods

The bivalves were collected from Abyan Coast (12°. 48. 485 N,

45°. 02. 381 E) in Aden Governorate. They were collected by hand during the spring low tides in the evening times to avoid higher temperatures and were then kept in open canvas sacs containing wet sand to minimize frictions, desiccation and then transported to the laboratory immediately. They were protected from agitation during the transportation. The clams were cleaned by gentle rubbing in clean seawater to remove the clogged sediment and mucus and kept in aquaria of uniform size, 40 cm long, 25 cm wide 20 cm height, each containing clean and filtered seawater. Clams of uniform size of (47 ± 1) mm long were used in the study to avoid susceptible size-based variations in response to the test chemicals. At the end of the acclimatization the experimental organisms for 4 days must be in excellent condition to tolerate the experimental conditions. There should be less than 2% mortality during acclimatization [7] Four clams from each of the control and from the exposed sets were transferred (five and ten days exposure to cured oil concentrations of 0.5, 0.1 and 5.1 ml/l into beakers of 500 ml capacity containing 400 ml filtered seawater. The water columns of the beakers were sealed with a 2.3 cm to 3 cm layer of inert liquid paraffin to prevent exchange of gas with atmosphere; following the method of [5,8].

The oxygen consumption was determined for two and six hours in dark chambers to minimize the external stimulations. Then the oxygen contents were determined by Winkler's method for the experimental jars, the initial water and the control water without clams at the end of the experiment. After each experiment, the soft tissues were removed cleaned and dried at 80°C for 24 hours, and dry weights were determined. Standard deviations were calculated based on 4 determinations in ml O<sub>2</sub> consumed per hour per gm dry weight.

A condition index relating dry tissue weight to shell length was calculated by:

$$\text{Condition index} = \frac{\text{Dry Weight (gm)}}{\text{Shell length (mm)}} \times 100$$

**Citation:** Dobian AS, Al-Hagibi HA. Toxicity of Mareb Crude Oil on Intertidal Clam *Tivela Ponderosa* and its Effect on Oxygen Consumption under Laboratory Conditions. Ann Clin Pharmacol Toxicol. 2019; 1(2): 1010.

**Copyright:** © 2019 Aziz S. Dobian

**Publisher Name:** Medtext Publications LLC

**Manuscript compiled:** August 07<sup>th</sup>, 2019

**\*Corresponding author:** Hagib A. Al-Hagibi, Department of Earth and Environmental Sciences, Sana'a University, Yemen, E-mail: hagib\_2002@hotmail.com

The soft tissues were shucked off the shells, weighted then dried as above and reweighted to get the dry tissue weights. Then the dry-wet tissue weight ratio in percentage was calculated.

$$\text{Dry - Wet weight ratio} = \frac{\text{Dry Weight}}{\text{Wet weight}} \times 100$$

## Results

### Rate of oxygen consumption


Table 1 and Figure 1 illustrate the results of the rate of oxygen consumption of the control and Mareb Crude Oil exposed clams ml O<sub>2</sub>/g dry soft tissues/hr for five and ten days (every two and six hours) of observation. Oxygen consumption in the control was 0.131 and 0.121 ml/g dry weight/hr. The oxygen uptake increased gradually in clams exposed to different concentrations of the crude oil, in the low concentration 0.5 ml/l was 0.135 and 0.124 ml/g dry weight/hr, and the medium concentration 0.1 ml/l was 0.141 and 0.137 ml/g dry weight/hr, and the increase reached its peak in the clams exposed to high crude oil concentration 1.5 ml/l with an increase was 0.319 and 0.247 ml/g dry weight/hr, compare control during two and six hours respectively.

The results of the rate of oxygen consumption of the control and test substance exposed clams ml O<sub>2</sub>/g dry soft tissues/hr for ten days (every two and six hours) of observation. Oxygen consumption rate increased during the ten days compared oxygen consumption rate for five days. In the control was 0.222 and 0.126 ml/g dry weight/hr. During the low, medium and high concentration was {(0.561) (0.432)}, {(0.946) (0.573)} and {(1.117) (0.632)} ml/g dry weight/hr, respectively (Table 2).

Figure 1 and 2 show clearly the difference between the two groups of clams after 5 and 10 days exposure to Mareb Crude Oil toxicity (after two and six hours). The group in the (Figure 1 and 2) illustrate the rates of oxygen consumed by the control and the exposed clams after they have been transferred to crude oil-free seawater. In the 10 days group the rate of oxygen uptake was generally less than the rate of oxygen consumption in the 5 days group.

**Table 1:** Average oxygen consumption (ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup>) in bivalves exposed to Mareb Crude Oil for five days (after 2 and 6 hours) values are mean ± for 2 determinations.

Period of exposure (after six hours) Mean ± S.D	Period of exposure (after two hours) Mean ± S.D	Exposed (to ml/l)
0.121 ± 0.015	0.131 ± 0.009	0.0 (control)
0.124 ± 0.070	0.135 ± 0.010	0.5
0.137 ± 0.014	0.141 ± 0.020	1.0
0.247 ± 0.003	0.319 ± 0.029	1.5



**Figure 1:** Oxygen consumption by *Tivela ponderosa* after 5 days exposure to Mareb Crude Oil.

Finally, comparison of the charts after 5 and 10 days show clearly the difference between the two groups of clams after 5 days and two groups of clams after ten days of exposure to substance tests toxicity (after two and six hours). The charts illustrate the rates of oxygen consumed by the control and the exposed clams after they have been transferred to chemicals tests-free seawater.

### Condition index


The observed values of condition index which is the ratio percentage between the dry weight (g) and shell length (mm) are given in (Table 3 and Figure 3) which indicate a reduction in condition index of the clams exposed to Mareb Crude Oil for period of 5 and 10 days. The condition index of the control clams was 2.927 after 5 days and 2.751 after 10 days. Where decreasing condition index of the clams exposed to different concentrations Mareb Crude Oil during 5 and 10 days, in the low concentration 0.5 ml/l were 2.587 and 2.446 ml/l, while the medium concentration 0.1 ml/l were 2.452, and 2.238 ml/l, the high concentration 1.5 ml/l were 2.396 and 2.198 ml/l.

### Dry-Wet weight ratio

The data of the ratio between dry and wet weights of the control and exposed clams during 5 and 10 days are shown in Tables 4 and Figure 4. The decreased ratio was more or less inversely proportional to the degree for each test chemicals concentrations compared to the control. The ratio were 25.11 and 25.08 in the control and in the exposed clams to Crude oil compared to the control were {(24.34) (24.24), (22.89) (22.06) and (21.93) (21.12)}.

**Table 2:** Average oxygen consumption (ml O<sub>2</sub> gm<sup>-1</sup> dry tissue h<sup>-1</sup>) in bivalves exposed to Mareb Crude Oil for ten days (after 2 and 6 hours) values are mean ± for 2 determinations.

Period of exposure (after six hours) Mean ± S.D	Period of exposure (after two hours) Mean ± S.D	Exposed (to ml/l)
0.126 ± 0.000	0.222 ± 0.011	0.0 (control)
0.432 ± 0.000	0.561 ± 0.033	0.5
0.573 ± 0.005	0.946 ± 0.000	1.0
0.632 ± 0.000	1.117 ± 0.025	1.5





**Figure 2:** Oxygen consumption by *Tivela ponderosa* after 10 days exposure to Mareb Crude Oil.

**Table 3:** Condition index as a function of Mareb Crude Oil concentrations for five and ten days.

Condition index for <i>Tivela ponderosa</i>		
concentrations for ten days Mean ± S.D	concentrations for five days Mean ± S.D	Concentrations (ml/l)
2.751 ± 0.232	2.927 ± 0.134	0.0 (control)
2.446 ± 0.312	2.587 ± 0.326	0.5
2.238 ± 0.565	2.452 ± 0.289	1.0
2.198 ± 0.203	2.396 ± 0.182	1.5

**Table 4:** Dry-wet weight ratio as a function of Mareb Crude Oil concentrations five and ten days. Values are means  $\pm$  S.D, n =4.

Condition index for <i>Tivela ponderosa</i>		Concentrations (ml/l)
concentrations for ten days	concentrations for five days	
Mean $\pm$ S.D	Mean $\pm$ S.D	
25.08 $\pm$ 0.07	25.11 $\pm$ 0.44	0.0 (control)
24.24 $\pm$ 0.35	24.34 $\pm$ 0.36	0.5
22.06 $\pm$ 0.06	22.89 $\pm$ 0.06	1.0
21.12 $\pm$ 0.07	21.93 $\pm$ 0.44	1.5

**Figure 3:** Effect of Mareb Crude Oil concentrations on condition index of *Tivela ponderosa* after five and ten days exposure.**Figure 4:** Effect of Mareb Crude Oil concentrations on dry-wet weight ratio of *Tivela ponderosa* after five and ten days exposure.

## Discussion

### Rate of oxygen consumption

The concept of assessing ventilation rate and  $O_2$  consumption rate to explain toxicant stress arose out of the knowledge that metabolism and activity are interrelated. Variations in metabolic rate modify the scope for activity and the degree of activity affects metabolic rate. Usually, laboratory determinations of sub lethal stress can delineate linear or nonlinear responses. The respiratory rate of aerobic organisms represents the metabolic activity and thus the oxygen consumption is a useful measure of sub lethal effects on the animals which has been used to determine the extent of stress by various natural and man-made perturbations. The extent of modifications in oxygen consumption rate may be considered as great as pollutant-produced alterations [9].

The present study showed a significant increase in oxygen consumption in the Mareb Crude Oil treated clams over that of the control clams. Oxygen consumption increased gradually with the increasing crude oil concentrations. The increase in oxygen consumption in the treated clams can be explained by the high metabolic activity of the organisms due to the stress imposed by the pollutant. The metabolic functions demand energy and subsequently oxygen is needed for oxidation. It was observed by [10] that in animals exposed to seawater extracts of the oil, from which particulate oil had

been removed, the metabolic rate was significantly stimulated; the degree of stimulation increased with increasing oil concentration. This leads one to explain the increase in oxygen consumption on the basis of oxygen utilization by the organism as due to demand for its high metabolic rate, which involves two distinct metabolic components; the basal metabolism reflects routine maintenance processes of the organism at rest and a further significant fraction of the oxygen demand is attributed to locomotion and other forms of activity [11].

The decline in oxygen consumption in the clams exposed to test crude oil concentrations during 10 days may be due in part, to the narcotization effect of the chemical on the animals and to the disturbance of the overall vital functions of the organisms by the compound. This condition can be attributed to the suppression effect of crude oil concentrations on the clams in addition to the effect of long term exposure. On the other hand, the decrease in oxygen consumption in case of the high hydrocarbon concentration is the result of activity reduction as [10] hypothesized this situation by stating that, the decrease in oxygen consumption is the result of activity reduction which may mask a general increase in basic metabolic rate.

Struhsakes et al. [12] reported similar finding in fish exposed to benzene and showed that benzene penetrated readily into tissues and stimulated the respiration of fish; they opined that the increased respiration reflected a requirement for more oxygen to metabolize the benzene. Higher concentrations or extended exposure times resulted in a decline in metabolic rate and a possible narcotic effect arising from accumulation of benzene in the tissues.

Several workers reported the effects of hydrocarbons on the respiration of marine organisms but the result varies considerably and in some cases an increase of oxygen consumption is recorded while decrease in others. The variations in the consumption of oxygen may be due to the type of hydrocarbon or the methods of preparations are the duration of exposure or due to the environmental conditions of the experiments [13]. Avolizi and Nuwahyid [14], who recorded a depression in the respiratory rates of the bivalves, *Brachidontes* and *Donax* exposed to light crude oil which showed reversal conditions when exposed to high concentrations of the oil.

The clam *Anadara granosa* exposed to naphthalene for a short term (96 hrs) exhibited a reduction in its oxygen consumption [15]. On the other hand, several authors recorded observations similar to those of the present study where low hydrocarbon concentrations increases oxygen consumption and the high concentration decreased the respiratory rates. This is in accordance with [16,17] who recorded an increase in respiratory rates as a function of Water Soluble Fraction (WSF) of No. 2 Fuel. Gilfillan [18] reported an increase in metabolic rates when bivalves were exposed to low hydrocarbon concentrations and reduction of it when exposed to higher hydrocarbon concentrations [10,19,13].

The clam *Mya arenaria* had doubled its oxygen consumption when exposed to lower oil concentration and when they were exposed to greater oil concentration showed a depression in the oxygen consumption rate. Stainken and Neff [20,21] stated that the respiratory rate of early and late zoeae and megalops of mud crab were increased by exposure to phenanthrene and naphthalene in which the former had a greater effect than the latter on respiratory rate.

The above mentioned finding by different authors indicates that the metabolic response to hydrocarbons is more complex than the simple unidirectional inhibition or stimulation suggested by some studies [10].

## Condition index

This study has shown that both the condition index and the ratio of dry-wet weight tissues in *T. ponderosa* were significantly altered by the exposure to Mareb Crude Oil. The condition index parameter is another tool to interpret the growth rate of the animal and the actual energy balance indicating protein, carbohydrate and lipid catabolism to counteract the stressful conditions of pollution. The present results showed a gradual decrease of condition index as the test chemicals concentrations increased.

This condition was also observed by other authors like Roesijidi and Anderson in *Macoma balthica* and Stekoll et al. [22] in *Macomabalthica*. Both the studies related the decrease in condition index to the negative energy balance which indicates energy utilization rather than storage. Grauby and Spliid [23] recorded a highly significant negative correlation with PAH in the common mussel. According to Anderson [24] the time factor is necessary in obtaining a good result in condition index.

## Dry-Wet weight ratio

The dry-wet weight ration also showed significant variation from the control with the decrease in values corresponding to the increase in oil concentration. The faster decrease in the dry weight in relation to the wet tissue weight may be explained by the loss of dry weight for using their energy reserves due to oil exposure. This state probably occurred because of various factors such as decreasing in feeding accompanied by the increase in metabolic rate and reduction in filtering rates. The present results are in agreement with the conclusion recorded [25-30]. Who studied the clam *Meretrix casta* exposed to various Crude and fuel oils whereby they lost dry weight faster than wet weight. Stekoll et al. [22] reported similar results.

## Conclusion

The present study showed a significant increase in oxygen consumption in the test substances treated clams over that of the control clams. Oxygen consumption increased gradually with the increasing test substances concentrations. The increase in oxygen consumption in the treated clams can be explained by the high metabolic activity of the organisms due to the stress imposed by the pollutant. On the other hand, when increasing the exposure period for Pollution substance decrease in oxygen consumption in the clams exposed to toxic substance concentrations. The possible reason for the latter is the insensitivity of the clams especially in high concentration of pollution substances.

## References

- Sprague JB. Measurement of pollutant toxicity to fish- III. Sublethal effects and "safe" concentration. *Water Res.* 1971;5(6):245-66.
- Swedmark M, Breaten B, Emanuelsson E, Granmo A. Biological effects of surface active agents on marine animals. *Mar Biol.* 1971;9(3):183-201.
- Deshmukh RS. On the oxygen consumption of estuarine mollusc, *Meretrix meretrix* under various conditions. *J Mar Biol Ass.* 1979;1-9.
- Prabhudeva KN, Menon NR. Oxygen consumption under copper and zinc stress in *Perna viridis*. *Fish Technol.* 1986;23(1):24-26.
- Mohan CV, Gupta TRC, Shetty HPC, Menon NR. Combined toxicity of mercury and cadmium to the tropical green mussel *Perna viridis*. *Dis Aquat Org.* 1986;2:65-72.
- Mohan CV, Menon NR, Gupta TRC. Filtration in some tropical intertidal bi valves exposed to mercury and cadmium mixtures. *Fish Technol.* 1986;23(2):204-10.
- Rand MC, Greenberg AE, Taras MJ. Bioassay methods for aquatic organisms. In: Standard methods for the examination of water and wastewater. 14th ed. American Public Health Association; 1976.
- Mathew R, Menon NR. Oxygen consumption in tropical bivalves, *Perna viridis* and *Meretrix casta* exposed to heavy metal. *Indian J Mar Sci.* 1983;12(1):57-9.
- Anderson JW. Responses to sub lethal levels of petroleum hydrocarbons: Are they sensitive indicators and do they correlate with tissue contamination. In: *Fate and Effects of Petroleum Hydrocarbons in Marine Ecosystems and Organisms.* 1977;95-114.
- Percy JA. Effect of dispersed crude oil upon the respiratory metabolism of an Arctic marine amphipod. *Onisimus (Boekisimus) affinis*. In: *Fate and Effects of Petroleum Hydrocarbons in Marine Ecosystem and Organisms*, Pergamon Press. 1977;192-200.
- Newell RC. *Biology of Intertidal Animals.* Elsevier. 1970.
- Struhsakes JW, Maxwell BE, Echeverria T. Effects of benzene (a water-soluble component of crude oil) on eggs and larvae of Pacific herrings and northern anchovy. 1974;253-284.
- Thomas RE, Rice SD. The effect of exposure temperature on oxygen consumption and opercular breathing rates of pink salmon fry exposed to toluene, naphthalene and water-soluble fraction of Cook Inlet crude oil and No.2 fuel oil. In: *Marine Pollution: Functional Responses.* 1979;39-51.
- Avolizi RJ, Nuwahyid M. Effects of crude oil and dispersants on bivalves. *Mar. Poll Bull.* 1974;5(10):149-53.
- Eapen JT. Physiological and biochemical responses of an acrid blood clam to naphthalene. Unpublished Ph.D. Thesis. 1987.
- Anderson JW, Neff JM, Cox BA, Tatem HE, Hightower GM. The effect of oil on estuarine animals: Toxicity, uptake and depuration, respiration. In: *Pollution and physiology of marine Organism.* 1974;285-310.
- Anderson JW, Neff JM, Cox BA, Tatem HE, Hightower GM. Characteristic of dispersion and water-soluble extracts of crude and refined oils and their toxicity to estuarine crustaceans and fish. *Mar Biol.* 1974;27(1):75-88.
- Gilfillan ES. Decrease of net carbon flux in two species of mussels caused by extracts crude oil. *Mar Biol.* 1975;29(1):53-7.
- Tatem H. Accumulation of naphthalene by grass shrimp: Effects on respiration, hatching and larval growth. In: *Fates and Effects of petroleum hydrocarbons in marine Ecosystems and Organisms.* Bull. 1977;19(11):201-9.
- Stainken D. The accumulation and depuration of No. 2 fuel oil by the soft shell clam *Mya arenaria*. In: *Fat and Effects Petroleum Hydrocarbons in Marine Ecosystem and Organisms.* Wolfe D.A, eds. 1977;313-22.
- Neff JM. Polycyclic Aromatic Hydrocarbons in the Aquatic Environment: Sources, Fates and Biological Effects. 1979.
- Stekoll MS, Clement LE, Shaw DG. Sublethal effects of chronic oil exposure on the intertidal clam, *Macoma balthica*. *Mar. Biol.* 1980;57(1):51-60.
- Grauby K, Spliid NH. Hydrocarbons and organochlorines in common mussel from the Kattegat and Belts and their relation to condition indices. *Mar Poll Bull.* 1995;30(1):74-82.
- Anderson JW. An assessment of knowledge concerning the fate and effects of petroleum hydrocarbons in the marine environment. In: *Marine Pollution: Function Responses.* 1979:3-21.
- Sophia AJ, Balasubramanian T. Changes in the physical condition of *Meretrix casta* exposed to water soluble fraction of refined and crude oil. *Arch Environ Contam Toxicol.* 1992;22(4):471-74.
- Baby KV. Combined toxicity of petroleum hydrocarbons on selected Ph.D. Thesis. Cochin University Technology, Cochin. 144 pp. heavy metals and marine organisms. *Of Science and Technology, Cochin.* 144 pp. 1987.
- Bayne BL. *Marine Mussels; Their Ecology and physiology*, International Biological Programme-10, Cambridge University Press Cambridge. 1976;1-495.
- Casalderrey FA, Ferrando MD, Andreu-Moliner E. Effect of the insecticide methyl parathion on filtration and ingestion rates of *Brachionus calyciflorus* and *Daphnia magna*. *Sci Total Environ.* 1993;134:867-76.
- Fong WC. Uptake and retention of Kuwait crude oil and its effects on oxygen uptake by the soft clam, *Mya arenaria*. *J Fish Res Board Can.* 1976;33:2744-80.
- Menon NN. Investigations On The Effects Of Petroleum Hydrocarbons On The Marine Bivalves *Perna Viridis (Linnaeus)* And *Sunetta Scripta (Linne)*. 1997.